

Research Article

Open Access

Construction of Epitope-based Vaccine for Avian Reovirus

Running title: Epitopes-based vaccine for avian reovirus

Hongzhuan Wu^{1*}, Lula Smith¹, Karyn Scissum-Gunn², Joseph J Giambrone³, Boakai K Robertson¹ and Robert Villafane¹

¹Department of Biological Sciences, Alabama State University, Montgomery, AL, 36101, USA

²Associate Vice President for Academic Operations, California State University, Fullerton, CA, 92831, USA

³Department of Poultry Science, Auburn University, AL 36849, USA

Article Info

***Corresponding author:**

Hongzhuan Wu

Professor

Department of Biological Sciences

Alabama State University

Montgomery, AL, 36101

USA

E-mail: hwu@alasu.edu

Received: July 20, 2018

Accepted: September 5, 2018

Published: September 11, 2018

Citation: Wu H, Smith L, Scissum-Gunn K, Giambrone JJ, Robertson BK, Villafane R. Construction of Epitope-based Vaccine for Avian Reovirus. *Madridge J Clin Res*. 2018; 2(1): 50-54.

doi: 10.18689/mjcr-1000109

Copyright: © 2018 The Author(s). This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Published by Madridge Publishers

Abstract

Epitope-based vaccine prototypes can stimulate protective immune responses against viruses. Identification of protective epitopes by traditional methods such as phage display is time and labor intensive. Such methods rely upon activity of epitope regions through empirical means for identification. Bioinformatics enables researchers to apply predictive approaches to identify protective epitopes. Software programs can predict epitopes by computer-driven pattern matching algorithms or analysis of conserved protein sequences for homology with known epitopes, however, such database services are expensive and limited to mouse and human pathogens. Avian reovirus (ARV) σ C protein is the main immunogenic surface protein of ARV. In this study, we developed a protocol for epitope prediction for an avian virus. Our predictions are based upon: (1) 2D and 3D structural analysis, (2) PROSITE glycosylation patterns, (3) protein sequence homology and alignments, and (4) hydrophobic index. Using this approach, we predicted 3 regions within the σ C protein that harbor putative protective epitopes. Then, we expressed 10 different size of recombinant σ C protein in a yeast expression system; the subject cell-neutralization of these recombinant proteins with avian reovirus S1133 strain in chicken fibroblast cells (CEF) not exactly collaborate our prediction.

Keywords: Reovirus; Sigma C; Epitopes; Bioinformatics.

Introduction

Avian reovirus (ARV) and mammalian reovirus (MRV) belong to the genus Orthoreovirus. Both share physical-chemical and morphological characteristic, including segmented genomes consisting of 10 genome segments of double-stranded (ds) RNA. The RNA is packaged into a non-enveloped icosahedral double capsid [1]. The genomic segments can be separated by polyacrylamide gel electrophoresis (PAGE) into three size classes, L (large), M (medium), and S (small). However, ARV differs from its mammalian counterpart in its lack of haemagglutination activity [2], ability to induce cell fusion [3, 4], and association with naturally occurring pathological conditions [5]. ARV is an important cause of diseases in poultry. In particular, reovirus-induced arthritis, chronic respiratory diseases, and malabsorption syndrome [6-9], provoke considerable economic losses.




Figure 1. Primer design diagram for sigma C peptide library construction

All ARV-encoded proteins, including 10 structural and 4 nonstructural proteins, have been demonstrated [10, 11]. Protein sigma C, encoded by the S1 gene [12], is the target for type-specific neutralizing antibodies [13]. Recently, ARV sigma A encoded by the S2 gene [14] has been identified as a double-stranded RNA (dsRNA) binding protein [14, 15] and possible involvement in resistance to interferon [16]. Another protein of ARV, Sigma NS, encoded by the S4 gene [17], has been reported for its single-stranded RNA (ssRNA) binding activity [18]. To date, no investigations regarding the epitope mapping of anti-sigma C monoclonal antibodies have been reported.

Among mammalian reoviruses (MRVs), there are three structural proteins shown to carry neutralizing epitopes: $\sigma 1$, $\sigma 2$, and $\lambda 2$. Hayers *et al* showed that $\sigma 1$ protein carried type-specific neutralizing epitopes, whereas $\sigma 2$ and $\lambda 2$ carried group-specific neutralizing epitopes. ARV-encoded proteins, including at least 10 structural proteins (λA , λB , λC , μA , μB , μBC , μBN , σC , σA , and σB), and 4 nonstructural proteins (μNS , P10, P17, and σNS) have been demonstrated. Protein σC , encoded by the S1 gene, shows noticeably higher divergence than other σ -class proteins and is a cell attachment protein and apoptosis inducer. Protein σC is also the target for type-specific neutralizing antibodies while antibodies against σB are group-specific. ARV σA encoded by the S2 gene has been identified as a double-stranded RNA (dsRNA) binding protein and possible involvement in resistance to interferon. Among avian reoviruses, structure-based sequence alignment of the ARV sigma C and mammalian reovirus (MRV) type 3sigma 1 have indicated the presence of heptad repeats and a triple alpha-helical coiled-coil structure in N-terminal region. In addition, crystallographic studies found that the carboxy terminal globular domain of ARV sigma C has a similar overall topology to that of MRV type 3 sigma 1. It has been shown that MRV sigma 1 protein recognizes the receptor of M cells (α -2-3 linked sialic acid) that facilitates penetration of antigens into intestinal Peyer's patches. However, there is little information on the ARV σC epitopes involved in the stimulation of host protective immunity. Moreover, current Bio-software are mostly mammalian-based and not commercial available, limited further bioinformatic analysis to its epitope region. In this study we first reported prediction of potential epitope regions on the

sigma C protein of avian reovirus strain S1133 by bioinformatic analysis, *S.pombe*- expressed deletion fragments of sigma C was produced, and its neutralizing ability with polyclonal antibody against avian reovirus in chicken embryo cells was tested.




Figure 2. Amplification of ARV σC with RT-PCR
M: 1kb DNA ladder; 1-10 corresponding to fragment 1-10

Materials and Methods

Bioinformatics' analysis of sigma C protein

The following structure were compared:(1) 2D and 3D structural analysis, (2) PROSITE glycosylation patterns, (3) protein sequence homology and alignments, and (4) hydrophobic index. Using this approach, we predicted 3 regions within the σC protein that harbor putative protective epitopes. Briefly, nucleotide and deduced amino acid sequences as well as the possible secondary structure of sigma C proteins were aligned and analyzed with a DNASTAR software package (DNASTAR Inc., Madison, WI, USA).The nucleotide sequence (981bp) of the full-length sigma C-encoding gene among ARV

isolates were aligned, corresponding to residues 1-327 of the sigma C protein. The cDNA sequences and predicted amino acid sequences of the sigma C were used in paired identity analysis to determine the extent of nucleotide and amino acid sequence identity and divergence. The alignments were further analyzed with phylogeny program.




Figure 3. Predicted 2-D Structure of ARV sC Protein

Gene expression and protein purification

The sigma C open reading frame was amplified by reverse-transcriptase polymerase chain reaction (RT-PCR). Primers for amplification of the 10 putative epitope region of σ C were obtained from Invitrogen, Inc. Carlsbad, CA.

Primers used in this study:

Primer Description	Strand	F-Forward R-Reverse	Primer length (bp)	Sequences	Corresponding position in sigma C ORF	Amplified fragment length(bp)	Tm (°C)
ARV-01	F	16	TCCATCGCAGCGAAGA	15-54	39	48	
ARV-02	R	17	TGACAGTATACAAGCTG				
ARV-03	F	17	ATTGACTTCGAACGTGA	54-450	396	48	
ARV-04	R	18	TTTAACACGATCCTGCAG				
ARV-05	F	17	ACATAACGCATATCCA	750-816	66	48	
ARV-06	R	15	AACGACGCAGCTTGG				
ARV-07	F	17	ATCATTGGAGTCTACCG	450-752	302	48-49	
ARV-08	R	17	TGCTCACTTAAATCGAAGG				
ARV-09	F	17	AGAGGTGAGCTTGATAC	30-381	351	49	
ARV-10	R	15	GGCAGTGGAGTTTCC				
ARV-11	F	19	ATCTCCAATTGAAGAGTG	382-661	279	48-49	
ARV-12	R	16	ACAGTGAGCGTTAACG				
ARV-13	F	15	GAATCCCGCACTGGG	662-802	240	50-52	
ARV-14	R	16	CATGGAAGACGCACTG				
ARV-15	F	25	ATGGCGGGTCTCAATCCATCGCAGC	1-981	981	67	
ARV-16	R	30	CTTATCATCATCATCGGTGTCGATGCCGT				
ARV-15+ARV-2	F+R	25+17		1-54	54	54	
ARV-13+ARV-16	F+R	15+30		662-981	319	52	

Table 1. Predicted protein molecular weight (MW)

Fragment & Position	Predicted MW(Dalton)
(1) 15-54	1597.84
(2) 54-450	13057.41
(3) 750-816	2423.71
(4) 450-780	11841.19
(5) 1-981	34855.76
(6) 30-381	12550.90
(7) 382-661	10064.20
(8) 662-802	5257.97
(9) 802-981	6424.97
(10) 1-15	504.61

Viral RNA was extracted from the S1133 strain using a Trizol kit (Invitrogen, Carlsbad, CA). Synthesis of cDNA was performed as described (). The PCR product was electrophoresis in a 1% agarose gel and visualized by ethidium-bromide staining. The PCR product was purified using a Gene clean II kit (Qbiogene, Inc., Carlsbad, CA) and inserted into the 6.0 kb yeast expression vector, pNMT1-TOPO (Invitrogen, Carlsbad, CA). The ten pNMT1- σ C constructs was mobilized into competent yeast (*Schizosachromycespombe*) cells (Invitrogen, Carlsbad, CA), and transformed yeast cells were selected on plates containing Thiamine on EMM medium (Invitrogen, Carlsbad, CA).

PROSITE Pattern	Positions on your sequence
ASN_GLYCOSYLATION, PATTERN. N-(P)(ST)(P)	27, 63, 102, 108, 127,
MYRISTYL, PATTERN. G-(EDRKHFFYW)-x(2)-[STAGCN]-[P]	8, 86, 128, 144, 271, 296, 311,
CK2_PHOSPHO_SITE, PATTERN. [ST]-x(2)-[DE]	29, 41, 53, 74, 122, 170, 308, 323,
LEUCINE_ZIPPER, PATTERN. L-x(6)-L-x(6)-L-x(6)-L	103,
PKC_PHOSPHO_SITE, PATTERN. [ST]-x-[RK]	9, 106, 185, 223, 297, 319,

Figure 4. Protein Patterns Search of ARV σ C

The ten different size of sigma C proteins was expressed in an *S. Pombe* expression system (Invitrogen, Carlsbad, CA) and purified by Pro Bond purification system (Invitrogen, Carlsbad, CA) according to manufacturer's instructions. Briefly, transformed yeast containing thepNMT1- σ C or the control plasmid, pNMT1-TOPO without insert was used for further study. A single recombinant yeast colony containing the pNMT1- σ C construct was inoculated into 50 ml of EMM plus thiamine medium and grown overnight at 30° C with shaking. Cells were harvested by centrifugation at 1500x g for 5 min at room temperature. The supernatant was discarded. Cells were suspended in 50ml of EMM centrifuged, and the process was repeated. After final wash, cells were suspended in 50ml of EMM, and 500 μ l aliquots of starter culture were inoculated into 100 ml of EMM and incubated at 30°C with shaking for 18 hr. Cells were harvested by centrifugation at 1500 x g for 5 min at 4°C and washed in 10ml of 1 X TE buffer containing 100mM NaCl. Cells were centrifuged at 1500 x g for 5 min at 4 °C and pellet suspended in 1 ml of 1 XTE + 100mM NaCl, and centrifuged for 2 min at top speed in the micro centrifuge. The supernatant was removed and 400 μ l of

acid-washed glass beads added. Cells were broken apart using a Bead Beater (Scientific Industry Co., Bohemia, NY) at maximum speed for 45 seconds in a bead beater. Tubes were placed on ice for 5 min and this procedure repeated 5 times. Cells were centrifuged in a micro centrifuge for 2 min at maximum speed, and the supernatant was transferred to afresh tube. Protein concentration was determined by the Bradford method. The supernatant was further purified by a purification column (Invitrogen, Carlsbad, CA), briefly, add 8ml of supernatant to the column, bind for 30 mins, centrifuge (800xg) 5 mins, aspirate supernatant, wash with 8ml native wash buffer (provided by Invitrogen, Carlsbad, CA), centrifuge (800xg) 5mins, aspirate supernatant, repeat this process three more times, clamp the column in a vertical position and snap off the cap on the lower end. Elute the protein with 8ml native elution buffer (provided by Invitrogen, Carlsbad, CA), collect 1 ml fractions and analysis with SDS-PAGE and Western blot.




Figure 5. Multiple Alignment of ARV σ C with MRV σ Sequences




Figure 6. Using the Kyte-Doolittle method of calculating hydrophilicity over a window length of 30

SDS-PAGE and western blotting

Western blot was performed as described previously. Proteins were separated by SDS-PAGE, transferred onto a nitrocellulose membrane. The membrane was blocked with casein, and subsequently incubated for 1 hr with the chicken anti-reovirus sera that was diluted 1:100 in phosphate-buffered saline (PBS), washed three times with PBS, and incubated with rabbit anti-chicken immunoglobulin G (IgG)

conjugated with horseradish peroxidase (Sigma, St. Louis, MO) diluted 1: 3000. After 1 hr of incubation at 37 °C, the membrane was washed, and stained with Diaminonenzidine (Sigma, St. Louis, MO).




Figure 7. The predicted 3D-structure of the sigma C trimer: Each color chain is a sigmaC monomer. Structure is predicted for the regions from 122 to 316. There is long helix from 124 to 153, which forms the stem region. A loop (154-164) and two anti-parallel beta-sheets regions (165-176, 179-193) for the neck region. The mushroom like head is form by residues 196 to 316. The model is built by homology modeling method using the template of the avian reovirus sigma C 117-326, (PDB 2VRS)

DNA sequencing and sequence analysis

To confirm the right clone of putative epitopes, 10 different fragments of the sigma C gene were cloned and sequenced.

MAGLNPSQRREVVSLILSLTSNVNISHGDLTPYERLTNLEASTELLHRSISDIS
TTVSNISANLQDMTHILDVDTANLDGLRTTIVTALQDSVSILSTNVTDLTNRSS
AHIAAILSSLQTTVDGNSTAINLSKSDISSNGLAITDLQDRVKSLESTASHGLSF
SPPLSVADGVVSLDMDPYFCSQRVSLTSYSAEAQLMQFRWMARGTNGSSDT
IDMTVNAHCHGRRTDYMMSSGTNLTVTSNVLLTFDLSDITHIPSDLARLVP
SAGFQAAFPVDVSFTRDSATHAYQAYGVYSSSRVFTITFTPTGDDGTANIRSL
TVRTGIDT

Figure 8. Predicted epitopes of sC protein: Bioinformatic analyses of ARV sC protein revealed putative epitopes located at 3 different regions: a. 5~ 18; b.28 ~150; c.250 ~ 272

Neutralization Assay

To detect the efficacy of different peptide against ARV, Neutralization assay was performed in Chicken Embryo Fibroblast cells (CEF). Briefly, CEFs were prepared from 9-day-old specific-pathogen-free (SPF) chicken embryonating eggs and grown in 96-well microplates. Chicken polyclonal antibody was used. Serial twofold dilutions were prepared of 150µl antibodies in 150µl minimum essential medium (MEM) containing antibiotics. Virus controls, negative serum control from uninfected chickens were also included. The S1133 virus dilution was added in 150µl amounts to all wells. The plates were incubated at 37°C in 5% CO₂ for 1 hr. then 200µl of the different mixtures of virus and polyclonal antibody or virus, eptitope and polyclonal antibody was inoculated into CEF monolayers. After 60-min incubation at 37°C, inoculate were removed and monolayers were re-incubated with 1% FCS MEM. The neutralizing capacity was demonstrated by absence of CPE at 48 hr post-infection.




Figure 9. Neutralization activity analysis of yeast expressed sigma C fragment #1 represents Negative control, # 12 represents Positive control, #2-11 represents fragment 1-10

Results and Discussion

Expression and identification of sigma C proteins

To create the full-length and deletion fragments of the sigma C gene of S1133 strain, PCR amplification using distinct primer pairs was carried out. Examination of all amplified PCR products following electrophoresis in agarose gels revealed the expected sizes. The resulting sigma C of the expression plasmid constructs was successfully over-expressed in *S. pombe*. Each plasmid DNA was introduced into *S. pombe*, and protein extracts of the induced recombinant cells were analyzed by 12% SDS-PAGE and could be visualized by Coomassie brilliant blue staining. Analysis of transformed and expression-induced yeast by SDS-PAGE revealed the expressed sigma C protein. The purified protein and protein extracts of the induced recombinant cells were then analyzed by Western blotting.

Virus-neutralization (VN)

Bioinformatics' analyses of ARV Sigma C protein revealed putative epitopes located at 3 different regions: a. 5 ~ 18; b. 28 ~ 150; c. 250 ~ 272. Neutralization activity analysis of yeast expressed sigma C fragment revealed fragment 3, 6, 7, has the highest neutralization activity, which is corresponding to the position of sigma C protein of 750-816; 382-661 and 662-802. The actually result doesn't match with the prediction, it indicates current bioinformatics analysis we used doesn't fit for sigma C protein. Although our lab has produced two monoclonal antibodies against avian reovirus strain S1133, but no MAb had neutralizing activity against the tested reovirus, therefore, we are using polyclonal antibody in neutralization assay.

Acknowledgements

This research was supported by the National Science Foundation's Alliances for Graduate Education and the Professoriate (AGEP) Program, Grant No. 1432991, and USDA-IFAFS grant (#00-52100-9705). We are grateful to Teresa Dormitorio and Chia-Chen Weng for their technical help.

References

- Spandidos DA, Graham AF. Complementation between temperature-sensitive and deletion mutants of reovirus. *Journal of Virology*. 1975; 16: 1444-1452.
- Glass SE, Naqi SA, Hall CF, Karr KM. Isolation and characterization of a virus associated with arthritis of chickens. *Avian Dis*. 1973; 17(2): 415-424. doi: 10.2307/1589226
- Bodelon G, Labrada L, Martinez-Costas J, Benavente J. The avian reovirus genome segment S1 is a functionally tricistronic gene that expresses one structural and two nonstructural proteins in infected cells. *Virology*. 2001; 290(2): 181-191. doi: 10.1006/viro.2001.1159
- Bodelon G, Labrada L, Martinez-Costas J, Benavente J. Modification of late membrane permeability in avian reovirus-infected cells: viroporin activity of the S1-encoded nonstructural p10 protein. *J. Biol. Chem*. 2002; 277(20): 17789-17796. doi: 10.1074/jbc.M202018200
- Robertson MD, Wilcox GE. Avian reovirus. *Veterinary Bulletin*. 1986; 56: 155-174.
- Wood GW, Nicholas RAJ, Hebert CN, Thornton DH. Serological comparisons of avian reoviruses. *Journal of Comparative Pathology*. 1980; 90(1): 29-38. doi: 10.1016/0021-9975(80)90025-0
- Rosenberger JK, Olson NO. Viral arthritis. In Calnek BW, Barnes HJ, Beard CW, McDougald LR, and Y. M. Saif (eds.). *Diseases of Poultry*. 10th ed. Iowa State University Press; Ames, IA. 1997; 711-718.
- Hieronymus DR, Villegas P, Kleven SH. Identification and serological differentiation of several reovirus strains isolated from chickens with suspected malabsorption syndrome. *Avian Dis*. 1983; 27(1): 246-54. doi: 10.2307/1590390
- Wickramasinghe R, Meanger J, Enriquez CE, Wilcox GE. Avian reovirus proteins associated with neutralization of virus infectivity. *Virology*. 1993; 194(2): 688-96. doi: 10.1006/viro.1993.1309
- Varela R, Benavente J. Protein coding assignment of avian reovirus strain S1133. *J. Virol*. 1994; 68(10): 6775-6777.
- Liu HJ, Lee LH, Hsu HW, Kuo LC, Liao MH. Molecular evolution of avian reovirus: evidence for genetic diversity and reassortment of the S-class genome segments and multiple cocirculating lineages. *Virology*. 2003; 314: 336-349. doi: 10.1016/S0042-6822(03)00415-X
- Shapouri MR, Kane M, Letarte M, Bergeron J, Arella M, Silim A. Cloning, sequencing and expression of the S1 gene of avian reovirus. *J Gen Virol*. 1995; 76(6): 1515-20. doi: 10.1099/0022-1317-76-6-1515
- Wickramasinghe R, Meanger J, Enriquez CE, Wilcox GE. Avian reovirus proteins associated with neutralization of virus infectivity. *Virology*. 1993; 194(2): 688-96. doi: 10.1006/viro.1993.1309
- Yin HS, H.K. Shieh, L.H. Lee. Characterization of the double stranded RNA genome segment S3 of avian reovirus. *Journal of Virological Methods*. 1997; 67(1): 93-101. doi: 10.1016/S0166-0934(97)00080-3
- Martinez-Costas J, Gonzalez-Lopez C, Vakharia VN, Benavente J. Possible Involvement of the Double-Standard RNA-Binding Core Protein α A in the Resistance of Avian Reovirus to Interferon. *J. Virol*. 2000; 74(3): 1124-1131. doi: 10.1128/JVI.74.3.1124-1131.2000
- Chiu CJ, Lee LH. Cloning and nucleotide sequencing of the S4 genome segment of avian reovirus S1133. *Arch. Virol*. 1997; 142(12): 2515-2520.
- Yin HS, Lee LH. Identification and characterization of RNA-binding activities of avian reovirus non-structural protein σ NS. *J. Gen. Virol*. 1998; 79: 1411-1413. doi: 10.1099/0022-1317-79-6-1411
- Huang PH, Li YJ, Su YP, Lee LH, Liu HJ. Epitope mapping and functional analysis of sigma A and sigma NS proteins of avian reovirus. *Virology*. 2005; 332 (2): 584-595. doi: 10.1016/j.virol.2004.12.005