

## 2nd International Conference on ge Pharma & Nutrition, Health and Aging

August 1-2, 2019 Valencia, Spain

## Observations on Nutritional Diet of a Transylvania Lactating Women Group with Potentially Impact of Early Life Nutrition and Long Term Effects

Ramona Suharoschi¹\*, Simona Codruta Heghes², Ioana Ecaterina Pralea², Laura Ioana Gavrilas³, Lorena Filip³, Adela Viviana Sitar Taut⁴, Angela Cozma⁴, Adriana Fodor⁵, Crina Muresan⁵ and Cristina Adela Iuqa².7

Laboratory of Molecular Nutrition and Proteomics, Institute of Life Sciences, Food Science Department, Faculty of Food Science and Technology,

University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Romania

<sup>2</sup>Department of Drug Analysis, Iuliu Hatieganu University of Medicine and Pharmacy, Romania

<sup>3</sup>Department of Bromatology, Iuliu Hatieganu University of Medicine and Pharmacy, Romania

<sup>4</sup>Medical Clinic IV, Iuliu Hatieganu University of Medicine and Pharmacy, Romania

<sup>5</sup>Diabetes and Metabolic Diseases, Iuliu Hatieganu University of Medicine and Pharmacy, Romania

<sup>6</sup>Food Engineering Department, University of Agricultural and Veterinary Medicine of Cluj-Napoca, Romania

Proteomic and Metabolomic Department, Iuliu Hatieganu University of Medicine and Pharmacy, Romania

**B**reast milk composition of the lactating women highly vary with their nutrition habits and thus vary many nutrients that are essential for growth and development and are secreted in breast milk. The aim of the present study was to evaluate the energy and nutrient intake from food of lactating mothers during a self-selected diet in a Romanian sample population from Transylvania.

**Methods:** A total of 33 lactating women living in Cluj county, from Transylvania were surveyed and average intake was recorded by 7-day diary and a general questionnaire, both designed by the research team. An experienced member of the research team trained every woman.

Results: The results highlighted that energy intake was significantly lower compared with the level of energy intake recommended for lactating women and the distribution of macronutrients was misbalanced in the study group. The results indicate that lactating mothers have an inadequate intake of energy (1879.00 kcal/day for mothers in G1 vs. 2744 kcal/day recommended and 1770.08 kcal/day for mothers in G2, G3, G4 vs. 2803 kcal/day recommended). Results from the current study and from similar studies in different countries which reflect that lactating mothers have a poor energy and nutrient balance can be attributed to low intakes of nutritious food categories such as fruits, vegetables, legumes, low- fat dairy and oily-fish and high intakes of saturated fat rich foods. With such inadequate food intake, mothers might be at risk of depleting their energy and nutrient reserves, therefore affecting breast milk composition with a negative health impact.

**Conclusions:** In the present study, lactating mothers did not meet dietary recommendations of energy intake and macronutrient, except for proteins. These results highlight the importance that women, during breastfeeding, should receive proper information from professionals in order to improve their overall nutrition.

## **Biography:**

Ramona Suharoschi has more than 20 years' experience in the field of applied nutrigenomics in food science, development and optimization of nutritional intervention of functional food products, food safety and food toxicology. She had completed PhD in the field of Medical Sciences (Vet Medicine) at the University of Agricultural Sciences and Veterinary Medicine of Cluj-Napoca, Romania, following a research stage at the University of Reno, Nevada USA and the University of Ulster, Coleraine, Northern Ireland with the research topic "In silico Studies of Prostate Cancer". She had published more than 100 papers in national and international journals of scientific flow.